


Sex Differences in Hip Muscle Strength in Patients with Femoroacetabular Impingement Syndrome

Davis J^a, Brown L^b, Ellis T^{c,d}, Di Stasi S^{e,f}

^aExercise Science Program, College of Education and Human Ecology, The Ohio State University, ^bSchool of Health and Rehabilitation Sciences, The Ohio State University, ^cOrthopedic One, Ohio Orthopedic Surgery Institute, ^dDublin Methodist Hospital, ^eSports Medicine Research Institute, The Ohio State University Wexner Medical Center, ^fDepartment of Orthopaedics, The Ohio State University

Background

- Femoroacetabular impingement syndrome (FAIS) is a debilitating musculoskeletal condition that can cause hip pain and weakness, functional disability, and eventually osteoarthritis^{1,4}



Pictured above are the two types of bony lesions
Illustration by the American Orthopaedic Society for Sports Medicine

- Women with FAI have demonstrated poorer function both pre- and postoperatively compared to men;^{2,3,5} however, it is unclear what impairments may have contributed to these discrepancies.

Purpose: To investigate sex differences in hip strength for persons who have undergone hip arthroscopy for symptomatic FAIS. We tested the hypotheses that:

- Women would have significantly lower hip strength after surgery
- The involved limb would be weaker than the uninvolved limb, regardless of sex

Methods

Demographics | *Mean ± standard deviation*

	Males	Females	P =
N	12	13	-
Age (y)	28 ± 11	22 ± 11	0.225
BMI (kg/m ²)	25 ± 3	25 ± 5	0.774

Sports Participation Level at Testing


	Males (%)	Females (%)
Level 1	25	15
Level 2	25	0
Level 3	42	62
Level 4	8	23

Methods, Cont.

Prospective, Longitudinal Cohort Study

- Bilateral hip strength was assessed six months post arthroscopy
- Maximum voluntary isometric contraction (MVIC) was measured in side-lying position for:
 - Abduction
 - External rotation

Hip Strength Testing Positions




- Strength was normalized to body mass and limb averages were calculated

Statistical Analysis

- A two-way ANOVA with post-hoc t-tests were performed to investigate sex and limb differences in normalized strength (P<0.05)

Results



- There was no significant interaction of limb by sex (P=0.30) or significant effect of limb (P=0.347)
- There was a significant main effect of sex (P≤0.03) for hip abduction and external rotation strength
- Women demonstrated significantly lower normalized strength for both hip abduction and external rotation (P<0.002)

Results, Cont.

Strength by sex, regardless of limb | *Mean ± standard deviation*

Normalized strength represented as percent of body mass

	Males	Females	P =
Abduction	12.5 ± 3.3	9.4 ± 3.4	0.002
External Rotation	12.3 ± 3.1	9.0 ± 3.7	0.001

Discussion

- Our study supports our hypothesis that women have significantly lower hip strength than men after surgery.
- Strength deficits of the involved limb compared to the uninvolved limb were not identified in this cohort.
- These results may be due to a limitation of the methods used to assess muscle strength or the inclusion of only two major hip muscle groups.
- Other limitations include:
 - Small number of patients
 - Strength only measured in side-lying position
- Gaining a greater understanding of potential sex differences in strength will better equip clinicians to provide treatment to prevent functional deficits.
- Future studies should use larger groups of participants and evaluate other major muscle groups of the hip, which are commonly associated with FAI.

Acknowledgements

The project described was supported in part/full by the Legacy Fund of the Sports Physical Therapy Section of the American Physical Therapy Association.

References

- Casartelli N, et al. *Osteoarthritis and Cartilage*, 2011
- Joseph R, et al. *AJSM*, 2015
- Malviya A, et al. *The Bone & Joint Journal*, 2012
- Nepple J, et al. *Arthroscopy*, 2015
- Philippson M, et al. *Arthroscopy*, 2012

